

## Tilburg University

### A quadratically convergent parallel Jacobi process for almost diagonal matrices with distinct eigenvalues

Paardekooper, M.H.C.

*Publication date:*  
1987

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

Paardekooper, M. H. C. (1987). *A quadratically convergent parallel Jacobi process for almost diagonal matrices with distinct eigenvalues*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 289). Unknown Publisher.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM  
R

6

7626  
1987  
289

89

BRABANT

POSTBOX 90153  
5000 LE TILBURG  
THE NETHERLANDS




DEPARTMENT OF ECONOMICS  
RESEARCH MEMORANDUM

A QUADRATICALLY CONVERGENT PARALLEL  
JACOBI PROCESS FOR ALMOST DIAGONAL  
MATRICES WITH DISTINCT EIGENVALUES

M.H.C. Paardekooper *wx 512.0*

FEW 289

*R 512*



K.U.B.  
BIBLIOTHEEK  
TILBURG

A QUADRATICALLY CONVERGENT PARALLEL JACOBI PROCESS

FOR ALMOST DIAGONAL MATRICES WITH

DISTINCT EIGENVALUES

M.H.C. Paardekooper

Department of Econometrics

Tilburg University

Tilburg, The Netherlands

This research is part of the VF-program "Parallele Algoritmiëk", THD - WI - 08185 - 25, which has been approved by the Netherlands Ministry of Education and Sciences.

A Quadratically Convergent Parallel Jacobi Process for Almost Diagonal Matrices with Distinct Eigenvalues.

Abstract

This paper discusses a generalization of the Jacobi process (1846). In each step  $\frac{1}{2}n$  pairs of non-diagonal elements are annihilated in an arbitrary almost diagonal matrix. We prove that the recursively constructed sequence of matrices converges to a diagonal matrix. The quadratically convergent method is adapted to parallel computation on an array processor.

## 1. INTRODUCTION

In the wellknown Jacobi algorithm [5] for the diagonalization of a real symmetric matrix each of the successive orthogonal similarity transformations is designed to annihilate a symmetric pair of off-diagonal elements. The ultimately quadratic convergence of this algorithm has been investigated by several authors [6,7,9,12].

In this paper we describe a natural, despite new generalization of that Jacobi process for the diagonalization of a non-normal almost diagonal matrix  $A \in \mathbb{C}^{n \times n}$  with distinct eigenvalues. We assume  $n$  to be even. In each iteration

$$A^{(k+1)} = S_k^{-1} A^{(k)} S_k, \quad k \in \mathbb{Z}, \quad (1.1)$$

with  $A^{(0)} := A$ ,  $\frac{1}{2}n$  pairs of symmetrically placed off-diagonal elements are annihilated. The transformation matrices  $S_k$  are not unitary. Consequently monotonic decrease of the Frobenius norms of the non-diagonal parts of  $A^{(k)}$  can not be guaranteed despite the annihilation of elements.

In [9] the same lack of monotonicity caused genuine difficulties in the proof of the quadratic convergence of the Eberlein algorithm [3,8] for the algebraic eigenproblem.

Each  $S_k$  is a direct sum of  $\frac{1}{2}n$  unimodular shears  $T_{i,k}$ ,  $i = 1, \dots, n/2$ ;

$$T_{i,k} = \begin{bmatrix} p_{i,k} & q_{i,k} \\ t_{i,k} & s_{i,k} \end{bmatrix} \begin{matrix} \leftarrow \ell(i,k) \\ \leftarrow m(i,k) \end{matrix} \quad (1.2)$$

$$\begin{matrix} \uparrow & \uparrow \\ \ell(i,k) & m(i,k) \end{matrix}$$

They annihilate  $\frac{1}{2}n$  pairs of the transformed matrix. In case of hermitian  $A$  each  $T_{i,k}$  is a unitary  $2 \times 2$  block. Our strategy for the pivots  $(\ell(i,k), m(i,k))$  aims to annihilate each off diagonal element exactly once in  $n-1$  successive steps:

$$\{(\ell(i,k), m(i,k)) \mid i=1, \dots, n/2, k=0, \dots, n-2\} = \{(i,j) \mid 1 \leq i < j \leq n\}. \quad (1.3)$$

If the difference of  $A^{(k)} = D^{(k)} + E^{(k)}$  and its diagonal part  $D^{(k)} = \text{diag}(a_{jj}^{(k)})$  is small relatively the separation of the eigenvalues, the transformation matrix  $S_k$  almost equals the unit matrix. Then the previously annihilated elements can increase only slowly. That phenomenon gives rise to the quadratic convergence:

$$\|E^{(n-1)}\|_F \leq \text{constant} \|E^{(0)}\|_F^2. \quad (1.4)$$

The ideas of Brent and Luk [2] for the solution of the symmetric eigenproblem on a systolic array can be carried over to our method. Further may be remarked that our procedure can be used as final stage in Sameh's (slowly convergent) parallel normreducing eigenvalue algorithm [10], in order to obtain a quadratically convergent process.

The pivots of  $T_{ik}$  are  $(\ell(i,k), m(i,k))$ ,  $i = 1, \dots, n/2$ , with  $\ell(i,k) m(i,k)$  and  $\ell(i,k) < \ell(j,k)$  for  $i < j$ . (1.5)

The caterpillar permutation  $K_n$  [2] generates a pivot strategy that annihilates each off-diagonal element exactly once in a sweep of  $\frac{1}{2} n(n-1)$  shear transformations. These are performed in  $n-1$  steps. Permutation  $K_n$  can be read off from figure 1.




fig. 1 Caterpillar permutation  $K_n$ .

If  $b^{(0)} = (1, 2, \dots, n)^T$ , then

$$(K_n b^{(0)})_i = \begin{cases} i+2 & , \quad 2 \leq i = 2k \leq n-2 \\ i-2 & , \quad 5 \leq i = 2k-1 \leq n-1 \\ i-1 & , \quad i = 3, n \\ i & , \quad i = 1. \end{cases} \quad (1.6)$$

With  $b^{(k)} = K_n^k b^{(0)}$  we have



$$\{\ell(i,k), m(i,k)\} = \{b_{2i-1}^{(k)}, b_{2i}^{(k)}\}, \quad i = 1, \dots, \frac{1}{2}n. \quad (1.7)$$

Figure 2 gives the pivot pairs  $(\ell(i,k), m(i,k))$  in the caterpillar order for  $n = 6$

$\begin{array}{c} i \\ \backslash \\ k \end{array}$	1	2	3
0	(1,2)	(3,4)	(5,6)
1	(1,4)	(2,6)	(3,5)
2	(1,6)	(2,3)	(4,5)
3	(1,5)	(2,4)	(3,6)
4	(1,3)	(2,5)	(4,6)

fig. 2

For the proof of the quadratic convergence of the cyclic generalized Jacobi-process the rule (1.3) is essential.

For the description of the process we divide  $A^{(k)}$  into a diagonal and a nondiagonal part  $D^{(k)}$  and  $E^{(k)}$  respectively.

$$E^{(k)} = A^{(k)} - D^{(k)}, \quad d_{ij}^{(k)} = \begin{cases} a_{ij}^{(k)} & , i = j \\ 0 & , i \neq j. \end{cases} \quad (1.8)$$

We set

$$\epsilon_k = \|E^{(k)}\|_F \quad (1.9)$$

and

$$\vartheta_k = \min_{i \neq j} |a_{ii}^{(k)} - a_{jj}^{(k)}|. \quad (1.10)$$

The initial matrix  $A$  is almost diagonal namely

$$\epsilon_0 \leq \eta/10, \quad (1.11)$$

where  $\eta$  denotes the separation of the spectrum of A:

$$\eta = \min_{i \neq j} |\lambda_i - \lambda_j|.$$

Condition (1.11) implies [1] that each eigenvalue  $\lambda_i$  is affiliated to exactly one diagonal element, say  $a_{ii}$

$$|\lambda_i - a_{ii}| \leq \epsilon_0 \quad (1.12)$$

Consequently  $|a_{ii} - a_{jj}| \geq \eta - 2\epsilon_0 \geq 8\epsilon_0$ . Thus

$$\epsilon_0 \leq \eta/8. \quad (1.13)$$

The organisation of our paper is as follows. In section 2 we present the parallel annihilators together with their properties and effects during the first step. In section 3 we describe the consequences of the annihilations in a complete sweep. In that section the main result is attained:

if  $n \geq 7$  and  $\epsilon_0/\eta \leq (10n)^{-1}$  then  $\epsilon_{n-1}/\eta \leq 2n(\epsilon_0/\eta)^2 \leq \frac{1}{5} \epsilon_0/n$ .

Section 4 discusses conclusions and some unsolved problems.

## 2. PARALLEL ANNIHILATORS. THE FIRST STEP

Without loss of generality we may assume that

$$(\ell(i,0), m(i,0)) = (2i-1, 2i), \quad i = 1, \dots, \frac{1}{2}n$$

as in the caterpillar strategy. Now we consider the annihilation in the first step,  $k = 0$ , with the annihilators  $T_{i,0}$ . For reasons of simplicity the subscript  $k = 0$ , referring to the first step will be omitted in the formulae of this section.

The shear transformation  $T_i$  only affects the elements in the  $(2i-1)$ -th and the  $2i$ -th row and column.

The Jacobi-parameters of  $T_i$  are  $p_i, q_i, r_i, s_i$ :

$$T_i = \begin{bmatrix} p_i & q_i \\ r_i & s_i \end{bmatrix}, \quad p_i s_i - q_i r_i = 1, \quad i = 1, \dots, \frac{1}{2}n. \quad (2.1)$$

Let matrix  $A$  be partitioned as

$$A = \begin{bmatrix} A_{11} & A_{21} & \dots & A_{1, \frac{1}{2}n} \\ A_{21} & A_{22} & \dots & A_{2, \frac{1}{2}n} \\ \vdots & \vdots & & \vdots \\ A_{\frac{1}{2}n, 1} & A_{\frac{1}{2}n, 2} & \dots & A_{\frac{1}{2}n, \frac{1}{2}n} \end{bmatrix} = (A_{ij}), \quad (2.2)$$

where each  $2 \times 2$  submatrix  $A_{ij}$  is given by

$$A_{ij} = \begin{bmatrix} a_{2i-1, 2j-1} & a_{2i-1, 2j} \\ a_{2i, 2j-1} & a_{2i, 2j} \end{bmatrix}, \quad i, j = 1, \dots, \frac{1}{2}n. \quad (2.3)$$

For the  $2 \times 2$  blocks  $A_{ij}^{(1)}$  of  $A^{(1)} = S^{-1}AS$ , where  $S = \text{diag}(T_i)$ , we find

$$A_{ij}^{(1)} = T_i^{-1} \otimes T_j^T A_{ij}. \quad (2.4)$$

Especially for the diagonal block  $A_{ii}^{(1)}$ :

$$\begin{bmatrix} a_{2i-1,2i-1}^{(1)} \\ a_{2i-1,2i}^{(1)} \\ a_{2i,2i-1}^{(1)} \\ a_{2i,2i}^{(1)} \end{bmatrix} = \begin{bmatrix} s_i p_i & s_i r_i & -q_i p_i & -q_i r_i \\ s_i q_i & s_i^2 & -q_i^2 & -q_i s_i \\ -r_i p_i & -r_i^2 & p_i^2 & p_i r_i \\ -r_i q_i & -r_i s_i & p_i q_i & p_i s_i \end{bmatrix} \begin{bmatrix} a_{2i-1,2i-1} \\ a_{2i-1,2i} \\ a_{2i,2i-1} \\ a_{2i,2i} \end{bmatrix}. \quad (2.5)$$

In order to simplify the formulae we set

$$\sigma_i = a_{2i,2i-1}, \mu_i = a_{2i-1,2i}, \nu_i = a_{2i-1,2i-1} a_{2i,2i}, \quad i=1, \dots, \frac{1}{2}n. \quad (2.6)$$

The annihilation of both  $\sigma_i$  and  $\mu_i$  requires the solution of a quadratic equation with discriminant  $\nu_i^2 + 4\sigma_i\mu_i$ .

With (1.13) we get  $|2\sigma_i\mu_i| \leq \epsilon^2 \leq 9^2/64 \leq |\nu_i|^2/64$ . Thus

$$|1 + 4\sigma_i\mu_i \nu_i^{-2}| \geq 31/32. \quad (2.7)$$

Let be

$$F_i := (1 + 4\sigma_i\mu_i \nu_i^{-2})^{\frac{1}{2}}, \quad \text{Re}(F_i) > 0, \quad 1 \leq i \leq \frac{1}{2}n.$$

Lowerbound (2.7) enables the formulation of

THEOREM 2.1 The shears

$$T_i = \begin{bmatrix} p_i & q_i \\ r_i & s_i \end{bmatrix} \begin{bmatrix} (\frac{1}{2} + \frac{1}{2F_i})^{\frac{1}{2}} & \frac{-\mu_i}{\nu_i (F_i(1+F_i)/2)^{\frac{1}{2}}} \\ \frac{\sigma_i}{\nu_i (F_i(1+F_i)/2)^{\frac{1}{2}}} & (\frac{1}{2} + \frac{1}{2F_i})^{\frac{1}{2}} \end{bmatrix}, \quad i=1, \dots, n/2 \quad (2.8)$$

annihilate both  $\sigma_i$  and  $\mu_i$ . Such a shear  $T_i$  is called the annihilator, related to the pivots in question.

PROOF. With (2.11) we derive that  $a_{2i,2i-1}^{(1)} = 0$  and  $a_{2i-1,2i}^{(1)} = 0$  if

$$\begin{bmatrix} p_i \\ q_i \end{bmatrix} = \alpha_i \begin{bmatrix} (1+F_i)/2 \\ \sigma_i \nu_i^{-1} \end{bmatrix}, \quad \begin{bmatrix} r_i \\ s_i \end{bmatrix} = \beta_i \begin{bmatrix} -\mu_i \bar{\nu}_i^{-1} \\ (1+F_i)/2 \end{bmatrix}, \quad \alpha_i, \beta_i \in \mathbb{C} \setminus \{0\}.$$

The condition  $p_i s_i - q_i r_i = 1$  implies  $\alpha_i \beta_i = 2(F_i + F_i^2)^{-1}$ . With  $\alpha_i = \beta_i = (2/(F_i + F_i^2))^{\frac{1}{2}}$  we get (2.15).  $\square$

In the sequel we need the estimates in

THEOREM 2.2. If  $\epsilon \leq \eta/10$  then the following inequalities hold for  $T_i$ ,  $i = 1, \dots, \frac{1}{2}n$ .

$$(a) \quad |p_i|^2 \leq 1 + (\epsilon/\eta)^2 \leq 1 + (\epsilon/\vartheta)^2; \quad (2.9)$$

$$(b) \quad |F_i + F_i^2|^{-1} \leq \frac{1}{2} + \frac{5}{4} \epsilon^2/\eta^2; \quad (2.10)$$

$$(c) \quad \|T_i\|_2^2 \leq 1 + 2 \epsilon/\eta \leq \frac{6}{5}. \quad (2.11)$$

$$(d) \quad |(F_i + F_i^2)/2|^{-\frac{1}{2}} \leq 1 + \frac{6}{5} \epsilon^2/\eta^2. \quad (2.12)$$

$$\begin{aligned} \text{PROOF (a). } |p_i|^2 &= \left| \frac{1}{2} + \frac{1}{2} (1+4 \sigma_i \nu_i^{-2})^{-\frac{1}{2}} \right| \leq \frac{1}{2} + \frac{1}{2} (1-2 \epsilon^2/\vartheta^2)^{-\frac{1}{2}}/\eta^{-2} \\ &\leq \frac{1}{2} + \frac{1}{2} (1-2(\epsilon/\eta)^2(1-2\epsilon/\eta)^{-2})^{-\frac{1}{2}} \end{aligned}$$

since  $\vartheta \geq \eta-2\epsilon$ . If  $|x| < 1/10$  then  $(1-2x^2(1-2x)^{-2})^{-\frac{1}{2}} \leq 1 + 2x^2$ .

$$\text{Hence } |p_i|^2 \leq 1 + \epsilon^2/\eta^2 \leq 1 + \epsilon^2/\vartheta^2.$$

$$\begin{aligned} (b) \quad |F_i + F_i^2|^{-1} &= |1+4 \sigma_i \mu_i \nu_i^{-2} + (1+4 \sigma_i \mu_i \nu_i^{-2})^{\frac{1}{2}}|^{-1} \\ &\leq (1-2x^2(1-2x)^{-2} + (1-2x^2(1-2x)^{-2})^{\frac{1}{2}})^{-1} \end{aligned}$$

where as above  $x = \epsilon/\eta$ . Simple but tedious calculations affirm

$$(1-2x^2(1-2x)^{-2} + (1-2x^2)(1-2x)^{-2})^{\frac{1}{2}}^{-1} \leq \frac{1}{2} + \frac{5}{4} x^2, \quad |x| \leq 1/10.$$

(c) Let be  $x_i = |p_i|^2 + |q_i|^2$ ,  $y_i = |r_i|^2 + |s_i|^2$ ,  $z = p\bar{r}_i + q\bar{s}_i$ . Then

$$T_i T_i^* = \begin{bmatrix} x_i & z_i \\ \bar{z}_i & y_i \end{bmatrix}$$

and  $x_i y_i - |z_i|^2 = 1$  as follows from the unimodularity of annihilator  $T_i$ .

So  $x_i + y_i = ((x_i - y_i)^2 + 4|z_i|^2 + 4)^{\frac{1}{2}} \geq 2$  and

$$\|T_i\|_2^2 = (x_i + y_i + ((x_i - y_i)^2 - 4)^{\frac{1}{2}})/2 \quad (2.13)$$

With (2.8) we get

$$\begin{aligned} x_i + y_i - 2 &= 2\left(\left|\frac{1}{2} + \frac{1}{2} F_i^{-1}|-1\right) + 2(|\sigma_i|^2 + |\mu_i|^2)/|\nu_i^2(F_i + F_i^2)|\right. \\ &\leq |(1-F_i)/F_i| + 2\epsilon^2/|\nu_i^2(F_i + F_i^2)| \\ &\leq (4|\sigma_i \mu_i| + 2\epsilon^2)/|\nu_i^2(F_i + F_i^2)| \leq (2+5\epsilon^2/\eta^2)\epsilon^2/\theta^2 \end{aligned}$$

as follows from (2.10). Hence

$$\|T_i\|_F^2 = x_i + y_i \leq 2 + 2.05 \epsilon^2/\theta^2 \quad (2.14)$$

From (2.13) and (2.14) it follows with easy calculations that

$$\|T_i\|_2^2 \leq 1 + \frac{8}{5} \epsilon/\theta \leq 1 + \frac{8}{5} \epsilon \eta^{-1} (1-2\epsilon/\eta)^{-1} \leq 1 + 2 \epsilon/\eta \leq \frac{6}{5}$$

for  $\epsilon \leq \eta/10$ .

(d) The derivation of (2.12) is similar to that of (2.10).  $\square$

An upperbound for the growth of norm of the nondiagonal part after the first step is given in

THEOREM 2.3. If  $\epsilon < \eta/10$ , then

$$\epsilon_1 \leq (1+2\epsilon/\eta) \epsilon \leq \frac{6}{5} \epsilon. \quad (2.15)$$

PROOF. The  $2 \times 2$  blocks in the partitioned  $A^{(1)}$  are

$$A_{ij}^{(1)} = T_i^{-1} A_{ij} T_j, \quad i, j = 1, \dots, \frac{n}{2}.$$

Since  $a_{2i-1, 2i}^{(1)} = a_{2i, 2i-1}^{(1)} = 0$ ,  $i = 1, \dots, \frac{n}{2}$ , with (2.11) we find

$$\begin{aligned} \epsilon_1^2 &= \sum_{i \neq j} \|A_{ij}^{(1)}\|_F^2 \leq \sum_{i \neq j} \|T_i^{-1}\|_2^2 \|T_j\|_2^2 \|A_{ij}\|_F^2 \\ &\leq \sum_{i \neq j} \|T_i\|_2^2 \|T_j\|_2^2 \|A_{ij}\|_F^2 \leq (1+2\epsilon/\eta)^2 \epsilon^2. \quad \square \end{aligned}$$

According to (1.12)  $a_{ii}^{(0)}$  is affiliated with  $\lambda_i$  and

$$|a_{ii}^{(0)} - \lambda_i| \leq \epsilon_0.$$

This affiliation is not changed by the annihilating transformation. This will be proved in

THEOREM 2.4. If  $\epsilon < \eta/10$  then the affiliation of the eigenvalues remains unchanged by the annihilating transformation

$$A^{(1)} = S^{-1} A S$$

where  $S = \text{diag}(T_i)$  with  $T_i$  given by (2.8)

PROOF. From (2.5) and (2.8) one gets

$$\begin{aligned} |a_{2i-1, 2i-1}^{(1)} - a_{2i-1, 2i-1}| &= |2 \sigma_i \mu_i \nu_i^{-1(1+F_i)}|^{-1} \\ &\leq \epsilon^2 |\nu_i|^{-2} (1+(1-2\epsilon^2 |\nu_i|^{-2})^{\frac{1}{2}})^{-1} |\nu_i| \\ &\leq (\epsilon/9)^2 (1+(1-2(\epsilon/9)^2)^{\frac{1}{2}})^{-1} |\nu_i| \leq |\nu_i|/126, \end{aligned} \quad (2.16)$$

for  $\epsilon/9 \leq 1/8$ . A change of affiliation only can be an exchange of affiliation between two elements at say the  $i$ -th shear transformation. Assume  $a_{2i-1,2i-1}$  and  $a_{2i-1,2i-1}^{(1)}$  to be affiliated with  $\lambda_{2i-1}$  and  $\lambda_{2i}$  respectively. Then as follows from (2.15)

$$|a_{2i-1,2i-1} - \lambda_{2i-1}| \leq \epsilon, \quad |a_{2i-1,2i-1}^{(1)} - \lambda_{2i}| \leq \frac{6}{5} \epsilon.$$

By (2.16)

$$\begin{aligned} |a_{2i-1,2i-1}^{(1)} - a_{2i-1,2i-1}| &\leq \frac{1}{126} |\nu_i| \\ &\leq \frac{1}{126} (|\lambda_{2i-1} - \lambda_{2i}| + 2\epsilon). \end{aligned} \quad (2.17)$$

The change of affiliation implies

$$|a_{2i-1,2i-1}^{(1)} - a_{2i-1,2i-1}| \geq |\lambda_{2i-1} - \lambda_{2i}| - \frac{11}{5} \epsilon. \quad (2.18)$$

Consequently we get with (2.17) and (2.18)

$$|\lambda_{2i-1} - \lambda_{2i}| - \frac{11}{5} \epsilon \leq \frac{1}{126} |\lambda_{2i-1} - \lambda_{2i}| + \frac{1}{63} \epsilon$$

Thus  $\epsilon > \frac{625}{1396} |\lambda_{2i-1} - \lambda_i| \geq \frac{2}{5} \eta$ . This contradicts (1.11).  $\square$



## 3. THE EFFECT OF A COMPLETE SWEEP

For the estimates of the norm  $\epsilon_k$  of the nondiagonal part  $E^{(k)}$  of  $A^{(k)}$ ,  $k = 0, \dots, n-1$ , we make use of the conclusion based on theorem 2.3: if  $\epsilon_k/\eta \leq \frac{1}{10}$  then

$$\epsilon_{k+1} \leq (1+2\epsilon_k/\eta)\epsilon_k. \quad (3.1)$$

In the first theorem we proof that a sufficient small  $\epsilon_0$  guarantees a slow growth of  $\epsilon_k$ .

THEOREM 3.1. If  $\epsilon_0 \leq \frac{1}{10} \eta n^{-1}$  then

$$\epsilon_k \leq \frac{1}{10} \eta (n-k)^{-1}, \quad k = 0, \dots, n-1. \quad (3.2)$$

PROOF. Assume  $\epsilon_k/\eta \leq \frac{1}{10} (\eta-k)^{-1}$ . Then, as follows from (3.1)

$$\epsilon_{k+1} \eta^{-1} \leq (1+2\epsilon_k \eta^{-1}) \epsilon_k \eta^{-1} \leq \frac{1}{10} (n-k)^{-1} (1 + \frac{1}{5}(n-k)^{-1}).$$

It is easy to verify that

$$\frac{1}{10} (n-k)^{-1} (1 + \frac{1}{5} (n-k)^{-1}) \leq \frac{1}{10} (n-k-1)^{-1}.$$

□

The transform  $A^{(k+1)}$  is obtained from  $A^{(k)}$  in two stages:

$$\begin{cases} A^{(k+\frac{1}{2})} = S_k^{-1} A^{(k)} = H^{(k)} A^{(k)} + B^{(k)}, \\ A^{(k+1)} = A^{(k+\frac{1}{2})} S_k = A^{(k+\frac{1}{2})} H^{(k)} + C^{(k)} \end{cases} \quad (3.3)$$

$$(3.4)$$

where  $H^{(k)}$  denotes the diagonal part of both  $S_k$  and  $S_k^{-1}$ .

Thus by (1.2)

$$\begin{cases} b_{\ell(i,k),j}^{(k)} = -q_{i,k} a_{m(i,k),j}^{(k)}, \quad b_{m(i,k),j}^{(k)} = -r_{i,k} a_{\ell(i,k),j}^{(k)}, \quad i = 1, \dots, n/2, \end{cases} \quad (3.5)$$

$$\begin{cases} c_{j,\ell(i,k)}^{(k)} = r_{i,k} a_{j,m(i,k)}^{(k+\frac{1}{2})}, \quad c_{j,m(i,k)}^{(k)} = q_{i,k} = q_{i,k} a_{j,\ell(i,k)}^{(k+\frac{1}{2})}, \quad j = 1, \dots, n. \end{cases} \quad (3.6)$$

Let be

$$G^{(k)} = B^{(k)} H^{(k)} + C^{(k)} . \quad (3.7)$$

Then

$$A^{(k+1)} = H^{(k)} A^{(k)} H^{(k)} + G^{(k)} . \quad (3.8)$$

According to (3.8) the updating  $A^{(k)} \rightarrow A^{(k+1)}$  comes about by two actions: a multiplication by two diagonal elements of  $S_k$  followed by the addition of  $g_{ij}^{(k)}$ . The multipliers  $H^{(k)}$  are rather tame as can be seen in

THEOREM 3.2. If  $\epsilon_0 \leq \frac{1}{10} \eta/n$  then

$$\prod_{k=2}^{n-2} \|H^{(k)}\|_2^2 \leq 155/154 .$$

PROOF. With (2.9) and (3.2), we get

$$\begin{aligned} \prod_{k=2}^{n-2} \|H^{(k)}\|_2^2 &\leq \prod_{k=2}^{n-2} (1 + \epsilon_k^2 / \eta^2) \leq \prod_{k=2}^{n-2} (1 + (n-k)^{-2} / 100) \\ &\leq \prod_{k=2}^{\infty} (1 + k^{-2} / 100) \leq 155/154 \end{aligned}$$

for  $\frac{1}{100} \sum_{k=2}^{\infty} k^{-2} = (\pi^2/6-1)/100$  and  $\exp (\pi^2/6-1)/100 \leq 155/154$  .  $\square$

The twofold action (3.8) for the construction of  $A^{(k+1)}$  is the starting point for the description on the history of the element  $a_{ij}^{(n-1)}$ ,  $i \neq j$ , after its annihilation in some step. Let the element in the position  $(i, j)$  be annihilated in step  $N(i, j) - 1 = N-1$ . Then

$$a_{i,j}^{(N)} = 0 , \quad a_{i,j}^{(N+1)} = g_{i,j}^{(N)} , \quad i \neq j . \quad (3.9)$$

and

$$a_{i,j}^{(k+1)} = h_{i,i}^{(k)} h_{j,j}^{(k)} a_{i,j}^{(k)} + g_{i,j}^{(k)} , \quad N+1 \leq k \leq n-2 . \quad (3.10)$$

So

$$|a_{i,j}^{(k+1)}| \leq \|H^{(k)}\|_2^2 |a_{i,j}^{(k)}| + |g_{i,j}^{(k)}|$$

and consequently

$$\begin{aligned} |a_{i,j}^{(n-1)}| &\leq \sum_{k=N}^{n-2} \prod_{m=k+1}^{n-2} \|H^{(m)}\|_2^2 |g_{i,j}^{(k)}| \\ &\leq \frac{155}{154} \sum_{k=N}^{n-2} |g_{i,j}^{(k)}|, \quad i \neq j, \end{aligned} \quad (3.11)$$

as follows from theorem 3.1.

The notational problems in the analysis of the parallel cyclic Jacobi-process are reduced with

DEFINITION 3.3. Let be  $V = (v_{ij}) \in \mathbb{C}^{n \times n}$ . Then  $\hat{V}^{(k)} = (\hat{v}_{ij}^{(k)})$  and  $\tilde{V}^{(k)} = (\tilde{v}_{ij}^{(k)})$ ,  $k = 0, \dots, n-2$ , are defined by

$$\hat{v}_{i,j}^{(k)} = \begin{cases} 0 & , i = j \text{ or } \{i,j\} \in \{\{\ell(i,k), m(i,k)\} \mid 0 \leq i \leq n/2\} \\ v_{i,j} & , (i,j) \text{ otherwise,} \end{cases} \quad (3.12)$$

$$\tilde{v}_{i,j}^{(k)} = \begin{cases} v_{i,j} & , i \neq j \wedge k > N(i,j) \\ 0 & , \text{ otherwise,} \end{cases} \quad (3.13)$$

$$\delta_k(V) := \|\hat{V}^{(k)}\|_F. \quad (3.14)$$

□

Figure 3 illustrates the pattern of zeros in the successive matrices  $\tilde{G}^{(k)}$ ,  $k=1, \dots, n-2=4$ . The  $\square$  indicates the position of an element annihilated in a preceding step. The corresponding elements in  $\tilde{G}^{(k)}$  contribute according (3.8) to  $a_{ij}^{(n-1)}$ .

$$\tilde{G}^{(1)} = \begin{bmatrix} 0 & \square & 0 & 0 & 0 & 0 \\ \square & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \square & 0 & 0 \\ 0 & 0 & \square & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \square \\ 0 & 0 & 0 & 0 & \square & 0 \end{bmatrix}; \tilde{G}^{(2)} = \begin{bmatrix} 0 & \square & 0 & \square & 0 & 0 \\ \square & 0 & 0 & 0 & 0 & \square \\ 0 & 0 & 0 & \square & \square & 0 \\ \square & 0 & \square & 0 & 0 & 0 \\ 0 & 0 & \square & 0 & 0 & \square \\ 0 & \square & 0 & 0 & \square & 0 \end{bmatrix};$$

$$\tilde{G}^{(3)} = \begin{bmatrix} 0 & \square & 0 & \square & 0 & \square \\ \square & 0 & \square & 0 & 0 & \square \\ 0 & \square & 0 & \square & \square & 0 \\ \square & 0 & \square & 0 & \square & 0 \\ 0 & 0 & \square & \square & 0 & \square \\ \square & \square & 0 & \square & 0 & 0 \end{bmatrix}; \tilde{G}^{(4)} = \begin{bmatrix} 0 & \square & \square & \square & 0 & \square \\ \square & 0 & \square & 0 & \square & \square \\ \square & \square & 0 & \square & \square & 0 \\ 0 & \square & \square & \square & 0 & \square \\ 0 & \square & \square & \square & 0 & \square \\ \square & \square & 0 & \square & \square & 0 \end{bmatrix}$$

Fig. 3. The matrices  $\tilde{G}^{(k)}$ ,  $k=1, \dots, n-2 = 4$ , in case of the caterpillar cyclic order.

Evidently  $|\tilde{G}^{(k)}| \leq |\hat{G}^{(k)}|$ . Our notation allows to reformulate (3.11) in matricial form:

$$|E^{(n-1)}| \leq \frac{155}{154} \sum_{k=1}^{n-2} |\tilde{G}^{(k)}| \leq \frac{155}{154} \sum_{k=1}^{n-2} |\hat{G}^{(k)}|.$$

Thus

$$\epsilon_{n-1} \leq \frac{155}{154} \sum_{k=1}^{n-2} \|\hat{G}^{(k)}\|_F. \quad (3.15)$$

In the proof of the quadratic convergence an essential role has been reserved for

THEOREM 3.4. Let be  $G^{(k)}$  as defined in (3.7). If  $\epsilon_0/\eta \leq (10n)^{-1}$ , then

$$\|\hat{G}^{(k)}\|_F \leq \frac{84}{83} \left(1 + \frac{7}{10} (n-k)^{-1}\right) \epsilon_k^2/\eta \quad (3.16)$$

PROOF. Remark that  $\|\hat{G}^{(k)}\|_F \leq \|P^{(k)}\|_2 \|\hat{B}^{(k)}\|_F + \|\hat{C}^{(k)}\|_F$ . Firstly we estimate  $\|\hat{C}^{(k)}\|_F$ . By (3.6) and (3.12)

$$\begin{aligned} \|\hat{C}^{(k)}\|_F &\leq \sum_{i=1}^{n/2} \left( \left( \sum_{j=1}^{n(i)} |c_{j,\ell(i,k)}^{(k)}|^2 \right)^{\frac{1}{2}} + \left( \sum_{j=1}^{n(i)} |c_{j,m(i,k)}^{(k)}|^2 \right)^{\frac{1}{2}} \right) \\ &\leq \sum_{i=1}^{n/2} \left( |r_{i,k}| \left( \sum_{j=1}^{n(i)} |a_{j,m(i,k)}^{(k+\frac{1}{2})}|^2 \right)^{\frac{1}{2}} + |q_{i,k}| \left( \sum_{j=1}^{n(i)} |a_{j,\ell(i,k)}^{(k+\frac{1}{2})}|^2 \right)^{\frac{1}{2}} \right). \end{aligned}$$

where  $\sum_{j=1}^{n(i)}$  denotes summation for all  $j$  except  $j = \ell(i,k), m(i,k)$ .

With (2.8) it follows that

$$\|\hat{C}^{(k)}\|_F \leq \sum_{i=1}^{n/2} \rho_{i,k} \left( |\sigma_{i,k}| \left( \sum_{j=1}^{n(i)} |a_{j,m(i,k)}^{(k+\frac{1}{2})}|^2 \right)^{\frac{1}{2}} + |\mu_{i,k}| \left( \sum_{j=1}^{n(i)} |a_{j,\ell(i,k)}^{(k+\frac{1}{2})}|^2 \right)^{\frac{1}{2}} \right)$$

where  $\sigma_{i,k} = a_{m(i,k),\ell(i,k)}^{(k)}$ ,  $\mu_{i,k} = a_{\ell(i,k),m(i,k)}^{(k)}$  and

$$\rho_{i,k} = |\nu_{i,k}^2 (F_{i,k} + F_{i,k}^2)^2|^{-\frac{1}{2}}.$$

Now  $|\nu_{i,k}|^{-1} \leq \eta^{-1} (1 - 2\epsilon_k/\eta)^{-1} \leq \eta^{-1} (1 - (n-k)^{-1}/5)^{-1} \leq (1 + (n-k)^{-1}/4)/\eta$  for

$n - k \geq 2$ . (2.12) and (3.2) imply  $|(F_{i,k} + F_{i,k}^2)/2|^{-\frac{1}{2}} \leq 1 + \frac{6}{5} \epsilon_k^2/\eta^2 \leq 1.012$ .

Hence

$$\begin{aligned} \|\hat{C}^{(k)}\|_F &\leq 0.506 (1 + (n-k)^{-1}/4) \left( \delta_k^2 (A^{(k+\frac{1}{2})}) + \sum_{i=2}^{n/2} (|\sigma_{ik}|^2 + |\mu_{ik}|^2)/\eta \right) \\ &\leq \frac{42}{83} (1 + (n-k)^{-1}/4) \|S_k\|_2^2 \epsilon_k^2/\eta \leq \frac{42}{83} (1 + (n-k)^{-1}/4) (1 + (n-k)^{-1}/5) \epsilon_k^2/\eta \end{aligned}$$

as follows from (2.11) and (3.2). Thus

$$\|\hat{C}^{(k)}\|_F \leq \frac{42}{83} (1 + (n-k)^{-1}/2) \epsilon_k^2/\eta.$$

In a similar way one gets, now with (2.9):

$$\begin{aligned} \delta_k(B^{(k)}P^{(k)}) &\leq \|\hat{B}^{(k)}\|_F \|\mathcal{H}^{(k)}\|_2 \leq \frac{42}{83} (1+(n-k)^{-1}/4) (1+(n-k)^{-2}/100) \epsilon_k^2/\eta \\ &\leq \frac{42}{83} (1+3(n-k)^{-1}/8) \epsilon_k^2/\eta . \end{aligned}$$

Hence

$$\|\hat{G}^{(k)}\|_F \leq \frac{84}{83} (1+7(n-k)^{-1}/10) \epsilon_k^2/\eta . \quad \square$$

THEOREM 3.5. If  $\epsilon_0/\eta \leq (10n)^{-1}$  then

$$\epsilon_{n-1} \leq \frac{54}{53} \sum_{k=1}^{n-2} (1 + \frac{7}{10} (n-k)^{-1}) \prod_{j=0}^{k-1} (1+(n-j)^{-1}/5)^2 \epsilon_0^2/\eta . \quad (3.17)$$

PROOF. By (3.15) and (3.16)

$$\epsilon_{n-1} \leq \frac{155}{154} \frac{84}{83} \sum_{k=1}^{n-2} (1 + \frac{7}{10} (n-k)^{-1}) \epsilon_k^2/\eta . \quad (3.18)$$

According to theorem 3.1  $\epsilon_k/\eta \leq 1/10$  for  $k = 0, 1, \dots, n-1$  if  $\epsilon_0/\eta \leq \frac{1}{10} \eta^{-1}$ . Hence by (3.1) and (3.2),

$$\epsilon_{k+1} \leq (1+(n-k)^{-1}/5) \epsilon_k, \quad k = 0, \dots, n-2$$

Thus

$$\epsilon_k \leq \prod_{j=0}^{k-1} (1+(n-j)^{-1}/5) \epsilon_0, \quad k = 0, \dots, n-1 . \quad (3.19)$$

So (3.17) is a consequence of (3.18) and (3.19).  $\square$

LEMMA 3.6. Let be

$$t_n = \sum_{k=1}^{n-2} (1 + \frac{7}{10} (n-k)^{-1}) \prod_{j=0}^{k-1} (1+(n-j)^{-1}/5)^2$$

Then for  $n \geq 3$

$$t_n \leq 11n/6 + 1$$

PROOF. It is easy to see that  $t_n$  satisfies the recurrence relation

$$t_{n+1} = (1 + \frac{1}{5} (n+1)^{-1})^2 (t_n + 1 + 7n^{-1}/5).$$

With starting value  $t_3 = \frac{27}{20} (16/15)^2$  one verifies that

$t_n \leq \frac{11}{6}n + 1$ ,  $n = 3, 4, \dots, 9$ . Now assume  $t_n \leq 11n/6 + 1$ . Then

$$t_{n+1} \leq (1 + \frac{1}{5} (n+1)^{-1})^2 (\frac{11}{6}n + 2 + 7n^{-1}/5) \leq 11(n+1)/6 + 1$$

for  $n \geq 9$ . □

This lemma together with theorem 3.6 enables us to finish the proof of the quadratic convergence as formulated in

THEOREM 3.7. If  $\epsilon_0/\eta \leq (10n)^{-1}$  then

$$\epsilon_{n-1} \leq c_n \epsilon_0^2/\eta \tag{3.20}$$

where  $c_n = \frac{54}{53} (11n/6 + 1)$ . So

$$\epsilon_{k(n-1)} \leq (c_n/\eta)^{2^k-1} \epsilon_0^{2^k}.$$

PROOF. (3.17) and (3.18) give

$$\epsilon_{n-1} \leq \frac{54}{53} (11n/6 + 1) \epsilon_0^2/\eta < \frac{1}{5} \epsilon_0$$

since  $\epsilon_0/\eta \leq (10n)^{-1}$ . This guarantees the quadratic convergence. □



## 4. CONCLUSIONS

It goes without saying that our proof shares important features with the proof of the analogous problem for real symmetric matrices [6,7,11,12]. As in the paper of Ruhe [9] the complexity of the proof rises from the non-unitary transformations. These hinder the monotonic decrease of the norm of the non-diagonal parts  $E^{(k)}$  of the successive matrices  $A^{(k)}$ .

The proof is restricted to the case of almost diagonal matrices in the sense that  $\|E^{(0)}\|_F$  is small relatively  $\eta = \min_{i \neq j} |\lambda_i - \lambda_j|$ , the separation of  $A$ 's spectrum.

The method clarifies that in the final stage of a norm reducing eigenvalue algorithm the annihilation procedure suffices for the convergence of the process and can be executed in stead of the complicated and time consuming normreducing shear transforms in both the Eberlein as in the Sameh algorithm [3,8,14].

The long sequence of worst case estimates brings about that despite their stepwise sharpness even the final bound (3.20) underestimates the convergence quite considerably. The uniform lowerbound  $\eta(1-2\epsilon_k/\eta) \leq \theta_k$  for all  $|a_{ii}^{(k)} - a_{jj}^{(k)}|$  especially leads to a pessimistic result.

A numerical result illustrates that the condition  $\epsilon_0/\eta \leq (10n)^{-1}$  is far too strong as appeared in many test cases. The matrix is of order 6.  $A = D + E$  with

$$D = \text{diag}(1, 36, 29, 22, 8, 15), \quad e_{ij} = 2.8(j-i). \quad (4.1)$$



The eigenvalues of A are 3.416218, 9.792723, 15.704854, 21.295146, 27.207277 and 33.583782. So  $\eta = 5.590292$ .

k	$\epsilon_k$	k	$\epsilon_k$	k	$\epsilon_k$
0	<u>15.336223</u>	7	3.647763	14	$0.973630 \cdot 10^{-3}$
1	22.305149	8	2.061459	15	<u><math>0.624204 \cdot 10^{-4}</math></u>
2	19.654710	9	0.698883	16	$0.631612 \cdot 10^{-5}$
3	15.030077	10	<u>0.257238</u>	17	$0.816325 \cdot 10^{-7}$
4	11.300451	11	$0.907285 \cdot 10^{-1}$	18	$0.544994 \cdot 10^{-8}$
5	<u>5.509465</u>	12	$0.595277 \cdot 10^{-1}$	19	$0.476108 \cdot 10^{-10}$
6	4.941476	13	$0.470223 \cdot 10^{-2}$	20	<u><math>0.473872 \cdot 10^{-15}</math></u>

Fig. 4. The sequence  $\epsilon_k$  with caterpillar cyclic order.

Figure 4 illustrates that the convergence occurs despite the large  $\epsilon_0/\eta$ . There rest interesting researchproblems. Firstly how to modify the algorithm in case of multiple eigenvalues. The papers [4,13] give useful information for the structure of an almost diagonal matrix with multiple eigenvalues. It seems reasonable to skip the shear transform  $T_{ik}$  if the corresponding diagonal elements of  $A^{(k)}$  are affiliated with equal eigenvalues.

The criterion, based on  $\min_{i \neq j} |a_{i,i}^{(k)} - a_{j,j}^{(k)}|$  and  $\epsilon_k$ , for the transfer from the normreduction in the Sameh process to the annihilation procedure is an important question for future research [14].

## 5. REFERENCES

1. Bauer, F.L. and C.T. Fike, Norms and Exclusion Theorems, Num. Math., 2, 1960, pp. 137-141.
2. Brent, R.P. and F.L. Luk, The Solution of Singular-value and Eigenvalue Problems on Multiprocessor Arrays, SIAM J. Sci.Stat. Comput., 6, 1985, pp. 69-84.
3. Eberlein, P.J., A Jacobi-like Method for the Automatic Computation of Eigenvalues and Eigenvectors of an Arbitrary Matrix, J. Soc. Indust. Appl. Math., 10, 1962, pp. 74-88.
4. Fan, K. and A.J. Hoffman, Lower Bounds for the Rank and Location of the Eigenvalues of a Matrix. Nat. Bur. Stand. Appl. Math. Series 39, 1954, pp. 117-130.
5. Jacobi, C.G.J., Über ein leichtes Verfahren die in der Theorie der Seculär Störungen vorkommenden Gleichungen numerisch aufzulösen, Crelle's J., 30, pp. 51-96.
6. Kempen, H.P.M. van, On the Convergence of the Classical Jacobi Method for Real Symmetric Matrices with Nondistinct Eigenvalues, Num. Math., 2, 1966, pp. 11-18.
7. Kempen, H.P.M. van, On the Quadratic Convergence of the Serial Cyclic Jacobi Method, Num. Math., 2, 19-22.
8. Paardekooper, M.H.C., An Eigenvalue Algorithm Based on Normreducing Transformations, Technische Universiteit Eindhoven, Thesis, 1969.
9. Ruhe, A., On the Quadratic Convergence of a Generalization of the Jacobi Method for Arbitrary Matrices, BIT, 8, 1968, pp. 210-233.

10. Sameh, A.H., On Jacobi and Jacobi-like Algorithms for a Parallel Computer. *Math. Comp.*, 25, 1971, pp. 579-590.
11. Schönhage, A., On the Quadratic Convergence of the Jacobi Process, *Num. Math.*, 6, 1964, pp. 410-412.
12. Wilkinson, J.H., Note on the Quadratic Convergence of the Cyclic Jacobi Process, *Num. Math.*, 4, 1962, pp. 296-300.
13. Wilkinson, J.H., Almost Diagonal Matrices with Multiple or Close Eigenvalues, *Lin. Alg. Appl.*, 1, 1968, pp. 1-12.
14. Paardekooper, M.H.C., Sameh's parallel eigenvalue algorithm revisited, in *Algorithms and Applications of Vector and Parallel Computers*, (H.J.J. te Riele, T.H.J. Dekker and H.A. van der Vorst, Eds.), Elsevier Science Publishers, 1987, pp. 351-371.

## IN 1986 REEDS VERSCHENEN

- 202 J.H.F. Schilderlinck  
Interregional Structure of the European Community. Part III
- 203 Antoon van den Elzen and Dolf Talman  
A new strategy-adjustment process for computing a Nash equilibrium in a noncooperative more-person game
- 204 Jan Vingerhoets  
Fabrication of copper and copper semis in developing countries. A review of evidence and opportunities
- 205 R. Heuts, J. van Lieshout, K. Baken  
An inventory model: what is the influence of the shape of the lead time demand distribution?
- 206 A. van Soest, P. Kooreman  
A Microeconomic Analysis of Vacation Behavior
- 207 F. Boekema, A. Nagelkerke  
Labour Relations, Networks, Job-creation and Regional Development. A view to the consequences of technological change
- 208 R. Alessie, A. Kapteyn  
Habit Formation and Interdependent Preferences in the Almost Ideal Demand System
- 209 T. Wansbeek, A. Kapteyn  
Estimation of the error components model with incomplete panels
- 210 A.L. Hempenius  
The relation between dividends and profits
- 211 J. Kriens, J.Th. van Lieshout  
A generalisation and some properties of Markowitz' portfolio selection method
- 212 Jack P.C. Kleijnen and Charles R. Standridge  
Experimental design and regression analysis in simulation: an FMS case study
- 213 T.M. Doup, A.H. van den Elzen and A.J.J. Talman  
Simplicial algorithms for solving the non-linear complementarity problem on the simplotope
- 214 A.J.W. van de Gevel  
The theory of wage differentials: a correction
- 215 J.P.C. Kleijnen, W. van Groenendaal  
Regression analysis of factorial designs with sequential replication
- 216 T.E. Nijman and F.C. Palm  
Consistent estimation of rational expectations models

- 217 P.M. Kort  
The firm's investment policy under a concave adjustment cost function
- 218 J.P.C. Kleijnen  
Decision Support Systems (DSS), en de kleren van de keizer ...
- 219 T.M. Doup and A.J.J. Talman  
A continuous deformation algorithm on the product space of unit simplices
- 220 T.M. Doup and A.J.J. Talman  
The 2-ray algorithm for solving equilibrium problems on the unit simplex
- 221 Th. van de Klundert, P. Peters  
Price Inertia in a Macroeconomic Model of Monopolistic Competition
- 222 Christian Mulder  
Testing Korteweg's rational expectations model for a small open economy
- 223 A.C. Meijdam, J.E.J. Plasmans  
Maximum Likelihood Estimation of Econometric Models with Rational Expectations of Current Endogenous Variables
- 224 Arie Kapteyn, Peter Kooreman, Arthur van Soest  
Non-convex budget sets, institutional constraints and imposition of concavity in a flexible household labor supply model
- 225 R.J. de Groof  
Internationale coördinatie van economische politiek in een twee-regio-twee-sectoren model
- 226 Arthur van Soest, Peter Kooreman  
Comment on 'Microeconometric Demand Systems with Binding Non-Negativity Constraints: The Dual Approach'
- 227 A.J.J. Talman and Y. Yamamoto  
A globally convergent simplicial algorithm for stationary point problems on polytopes
- 228 Jack P.C. Kleijnen, Peter C.A. Karremans, Wim K. Oortwijn, Willem J.H. van Groenendaal  
Jackknifing estimated weighted least squares
- 229 A.H. van den Elzen and G. van der Laan  
A price adjustment for an economy with a block-diagonal pattern
- 230 M.H.C. Paardekooper  
Jacobi-type algorithms for eigenvalues on vector- and parallel computer
- 231 J.P.C. Kleijnen  
Analyzing simulation experiments with common random numbers



- 232 A.B.T.M. van Schaik, R.J. Mulder  
On Superimposed Recurrent Cycles
- 233 M.H.C. Paardekooper  
Sameh's parallel eigenvalue algorithm revisited
- 234 Pieter H.M. Ruys and Ton J.A. Storcken  
Preferences revealed by the choice of friends
- 235 C.J.J. Huys en E.N. Kertzman  
Effectieve belastingtarieven en kapitaalkosten
- 236 A.M.H. Gerards  
An extension of König's theorem to graphs with no odd- $K_4$
- 237 A.M.H. Gerards and A. Schrijver  
Signed Graphs - Regular Matroids - Grafts
- 238 Rob J.M. Alessie and Arie Kapteyn  
Consumption, Savings and Demography
- 239 A.J. van Reeken  
Begrippen rondom "kwaliteit"
- 240 Th.E. Nijman and F.C. Palmer  
Efficiency gains due to using missing data. Procedures in regression models
- 241 S.C.W. Eijffinger  
The determinants of the currencies within the European Monetary System

## IN 1987 REEDS VERSCHENEN

- 242 Gerard van den Berg  
Nonstationarity in job search theory
- 243 Annie Cuyt, Brigitte Verdonk  
Block-tridiagonal linear systems and branched continued fractions
- 244 J.C. de Vos, W. Vervaat  
Local Times of Bernoulli Walk
- 245 Arie Kapteyn, Peter Kooreman, Rob Willemse  
Some methodological issues in the implementation  
of subjective poverty definitions
- 246 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel  
Sampling for Quality Inspection and Correction: AOQL Performance  
Criteria
- 247 D.B.J. Schouten  
Algemene theorie van de internationale conjuncturele en structurele  
afhankelijkheden
- 248 F.C. Bussemaker, W.H. Haemers, J.J. Seidel, E. Spence  
On  $(v,k,\lambda)$  graphs and designs with trivial automorphism group
- 249 Peter M. Kort  
The Influence of a Stochastic Environment on the Firm's Optimal Dyna-  
mic Investment Policy
- 250 R.H.J.M. Gradus  
Preliminary version  
The reaction of the firm on governmental policy: a game-theoretical  
approach
- 251 J.G. de Gooijer, R.M.J. Heuts  
Higher order moments of bilinear time series processes with symmetri-  
cally distributed errors
- 252 P.H. Stevers, P.A.M. Versteijne  
Evaluatie van marketing-activiteiten
- 253 H.P.A. Mulders, A.J. van Reeken  
DATAAL - een hulpmiddel voor onderhoud van gegevensverzamelingen
- 254 P. Kooreman, A. Kapteyn  
On the identifiability of household production functions with joint  
products: A comment
- 255 B. van Riel  
Was er een profit-squeeze in de Nederlandse industrie?
- 256 R.P. Gilles  
Economies with coalitional structures and core-like equilibrium con-  
cepts

- 257 P.H.M. Ruys, G. van der Laan  
Computation of an industrial equilibrium
- 258 W.H. Haemers, A.E. Brouwer  
Association schemes
- 259 G.J.M. van den Boom  
Some modifications and applications of Rubinstein's perfect equilibrium model of bargaining
- 260 A.W.A. Boot, A.V. Thakor, G.F. Udell  
Competition, Risk Neutrality and Loan Commitments
- 261 A.W.A. Boot, A.V. Thakor, G.F. Udell  
Collateral and Borrower Risk
- 262 A. Kapteyn, I. Woittiez  
Preference Interdependence and Habit Formation in Family Labor Supply
- 263 B. Bettonvil  
A formal description of discrete event dynamic systems including perturbation analysis
- 264 Sylvester C.W. Eijffinger  
A monthly model for the monetary policy in the Netherlands
- 265 F. van der Ploeg, A.J. de Zeeuw  
Conflict over arms accumulation in market and command economies
- 266 F. van der Ploeg, A.J. de Zeeuw  
Perfect equilibrium in a model of competitive arms accumulation
- 267 Aart de Zeeuw  
Inflation and reputation: comment
- 268 A.J. de Zeeuw, F. van der Ploeg  
Difference games and policy evaluation: a conceptual framework
- 269 Frederick van der Ploeg  
Rationing in open economy and dynamic macroeconomics: a survey
- 270 G. van der Laan and A.J.J. Talman  
Computing economic equilibria by variable dimension algorithms: state of the art
- 271 C.A.J.M. Dirven and A.J.J. Talman  
A simplicial algorithm for finding equilibria in economies with linear production technologies
- 272 Th.E. Nijman and F.C. Palm  
Consistent estimation of regression models with incompletely observed exogenous variables
- 273 Th.E. Nijman and F.C. Palm  
Predictive accuracy gain from disaggregate sampling in arima - models



- 274 Raymond H.J.M. Gradus  
The net present value of governmental policy: a possible way to find the Stackelberg solutions
- 275 Jack P.C. Kleijnen  
A DSS for production planning: a case study including simulation and optimization
- 276 A.M.H. Gerards  
A short proof of Tutte's characterization of totally unimodular matrices
- 277 Th. van de Klundert and F. van der Ploeg  
Wage rigidity and capital mobility in an optimizing model of a small open economy
- 278 Peter M. Kort  
The net present value in dynamic models of the firm
- 279 Th. van de Klundert  
A Macroeconomic Two-Country Model with Price-Discriminating Monopolists
- 280 Arnoud Boot and Anjan V. Thakor  
Dynamic equilibrium in a competitive credit market: intertemporal contracting as insurance against rationing
- 281 Arnoud Boot and Anjan V. Thakor  
Appendix: "Dynamic equilibrium in a competitive credit market: intertemporal contracting as insurance against rationing"
- 282 Arnoud Boot, Anjan V. Thakor and Gregory F. Udell  
Credible commitments, contract enforcement problems and banks: intermediation as credibility assurance
- 283 Eduard Ponds  
Wage bargaining and business cycles a Goodwin-Nash model
- 284 Prof.Dr. hab. Stefan Mynarski  
The mechanism of restoring equilibrium and stability in polish market
- 285 P. Meulendijks  
An exercise in welfare economics (II)
- 286 S. Jørgensen, P.M. Kort, G.J.C.Th. van Schijndel  
Optimal investment, financing and dividends: a Stackelberg differential game
- 287 E. Nijssen, W. Reijnders  
Privatisering en commercialisering; een oriëntatie ten aanzien van verzelfstandiging
- 288 C.B. Mulder  
Inefficiency of automatically linking unemployment benefits to private sector wage rates

Bibliotheek K. U. Brabant



17 000 01059368 0